

IN THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Claims 1-26 (Cancelled)

27. (Currently Amended) A method for obtaining a collagen-based matrix from a submucosa tissue source, comprising:

treating the submucosa tissue source with a disinfecting agent to provide a disinfected submucosa tissue source; and

removing the collagen-based matrix from the disinfected submucosa tissue

source.~~sources.~~

28. (Original) The method of claim 27, wherein the submucosa tissue source is from an alimentary tract of a mammal.

29. (Original) The method of claim 28, wherein the mammal is a pig.

30. (Original) The method of claim 29, wherein the submucosa tissue source is from the small intestine of a pig.

31. (Original) The method of claim 27, wherein the disinfecting agent is an oxidizing agent.

32. (Original) The method of claim 27, wherein the disinfecting agent is a peroxy compound.

33. (Original) The method of claim 32, wherein the disinfecting agent is an organic peroxy compound.

34. (Original) The method of claim 33, wherein the disinfecting agent is a peracid.

35. (Original) The method of claim 34, wherein the peracid is selected from the group consisting of peracetic acid, perpropionic acid and perbenzoic acid.

36. (Original) The method of claim 35, wherein the peracid is peracetic acid.

37. (Original) The method of claim 34, wherein said treating includes treating the submucosa tissue source with a medium containing an alcohol and the peracid.

38. (Original) The method of claim 37, wherein the alcohol has one to about six carbon atoms.

39. (Original) The method of claim 38, wherein the alcohol is selected from the group consisting of ethanol, propanols, and butanols.

40. (Original) The method of claim 39, wherein the alcohol is ethanol.

41. (Original) The method of claim 40, wherein the medium is an aqueous ethanol solution containing from about 0.1% to about 0.3% by volume peracetic acid.

42. (Original) The method of claim 34, wherein said treating includes treating the submucosa tissue source with a medium containing the peracid and having a pH of about 2 to about 6.

43. (Original) The method of claim 42, wherein the medium has a pH of about 2 to about 4.

44. (Original) The method of claim 43, wherein the peracid is peracetic acid, and the medium contains about 0.1% to about 0.3% by volume of peracetic acid.

45. (Original) A method for obtaining a collagen-based matrix from a submucosa tissue source, comprising:

providing a submucosa tissue source which has been treated with a disinfecting agent;

and

removing the collagen-based matrix from said submucosa tissue source.

46. (Original) The method of claim 45, wherein said submucosa tissue source is from a small intestine.

47. (Original) The method of claim 46, wherein said disinfecting includes treating the submucosa tissue source with an oxidizing agent.

48. (Original) The method of claim 47, wherein said treating includes contacting the submucosa tissue source with an aqueous medium containing the oxidizing agent.

49. (Original) The method of claim 47, wherein said treating includes contacting the submucosa tissue source with an aqueous medium containing a peroxy compound.

50. (Original) The method of claim 49, wherein the peroxy compound is a peracid.

51. (Original) The method of claim 50, wherein the peracid is peracetic acid.

52. (Original) The method of claim 51, wherein the medium comprises an alcohol.

53. (Original) The method of claim 52, wherein the alcohol is ethanol.

54. (Original) The method of claim 51, wherein the small intestine is from a pig.

Claims 55-63. (Cancelled)